

образовательной среды в современных условиях [Текст] / А. А. Лобачев, И. А. Мамаева // Образовательная деятельность вуза в современных условиях : материалы международной научно-методической конференции. - Караваево : Костромская ГСХА, 2017. - С. 34.

6. Козленкова, Е. Н. Взаимодействие преподавателя и студента в информационно-коммуникационной предметной среде [Текст] / Е. Н. Козленкова // Современные проблемы информатизации профессионального образования : материалы Международной научно-практической интернет-конференции. - М. : МГАУ, 2012. - С. 29-34.

7. Alipichev, A. Improving the credibility of pedagogical diagnostics in E-Learning [Text] / A. Alipichev, L. Nazarova, M. Shingareva, A. Siman // CEUR Workshop Proceedings. - 2020. - Vol. 2861. - P. 203-209.

СЕКЦИЯ «ИНОСТРАННЫЙ ЯЗЫК В ПРОФЕССИОНАЛЬНОЙ СФЕРЕ И ЛИНГВОСТРАНОВЕДЕНИЕ»

УДК 631.8

IMPACT OF FERTILISERS ON SOIL PROPERTIES AND BIOMASS YIELD UNDER A LONG-TERM SWEET SORGHUM CROPPING SYSTEM

Abudujiaba Zunimaimaiti, Post-graduate student of the Department of Agriculture and Experimental Methods of Federal State Budgetary Educational Institution of Higher Education "Russian Timiryazev State Agrarian University", japparzunun@mail.ru

Sukhodolov I.A., Post-graduate student of the Department of Vegetable Growing of Federal State Budgetary Educational Institution of Higher Education "Russian Timiryazev State Agrarian University", gotem1996@mail.ru

Scientific advisor: *Mazirov Mikhail Arnoldovich*, Doctor of Biological Sciences, Professor of Institute of Agrobiotechnology of Federal State Budgetary Educational Institution of Higher Education "Russian Timiryazev State Agrarian University"

Abstract: A continuous long-term field experiment (2008–2018) was conducted in Xinjiang, north-western China, to assess the impact of farmyard manure (FYM) and inorganic fertilizers on the sustainable biomass yield of sweet sorghum cultivar (Xingaoliang No3) and soil chemical properties. Seven treatments, associated with nitrogen (N), phosphorus (P), potassium (K), FYM, and their different combination, were compared with the control plot (CK). As a result, the treatments NP, PK, NK, NPK and NPKM significantly increased the average biomass yields by 30-48% over CK. The 12 t/ha FYM per year with NPK (NPKM) increased both the yield and total soluble solids (TSS) by 48% and 7.9% respectively, while the 18 t/ha/year application rate of FYM had an adverse effect on yield. Stem TSS, soil available N and K for all treatments decreased while soil organic carbon, soil total salt and the available P for FYM applied treatments increased over the years. The soil pH stabilized at 7.8–8.2 at the end. In conclusion, the 12 t/ha/year of FYM is the most efficient rate for a single application or

incorporation with inorganic fertilizers. A more reasonable application rate of N and K fertilizer to increase the yield and irrigation rate to reduce soil salt needs for further investigation.

Key words: *soil nutrient; biological mass; Sorghum bicolor (L.) Moench; salinity; arid region*

Sweet sorghum (*Sorghum bicolor* (L.) Moench) is a common salt-tolerant grain sorghum crop suitable for arid land cultivation and shows several advantages, e.g., high biomass yield, rapid growth, wide adaptability to diverse climate and soil conditions, high resistance to drought and salinity, and good yield potential in marginal environments (Zhang et al. 2016). Therefore, sweet sorghum has been the most dominant forage crop cultivated in Xinjiang. Some experimental achievements conducted in this study area showed that the sweet sorghum cultivar Xingaoliang No. 3 was relatively more resistant to soil salinity and suitable for accumulating higher sugar content than other cultivars (Zaituniguli et al. 2012), and it has been cultivated for years with different fertilization. The long-term application of farmyard manure (FYM) with balanced nitrogen (N), phosphorus (P), and potassium (K) fertilizer promotes soil microbial activity, prevents soil acidification and enhances crop production (Zhai et al. 2011, Chen et al. 2018). However, little attention has been paid to the change of soil fertility, the trend of annual crop yield, and the sugar contents of sweet sorghum for long-timescale in this region. Therefore, evaluating the fertilization management for sustainable sorghum production with higher stem sugar and improving soil condition became necessary. Hence, a continuous fertilization experiment in combination with FYM was conducted under the intensive long-term monocropping sweet sorghum system to examine the dynamics of biomass yield, stem sugar content, soil condition and available nutrients.

Material and Methods

Experimental site description. The field experiment was conducted for the sweet sorghum Xingaoliang No3, the conventional early-maturing cultivar, continuously from 2008 to 2018 at the experimental station of Xinjiang Academy of Agricultural Sciences, located in Manasi county (44°31'N, 86°26'E, 470 m a.s.l.), Xinjiang Uyghur Autonomous Region, China. The soil texture can be classified (USDA 1987) as silty loam. The soil covers can be termed anthrosols according to the soil classification system (FAO 2006).

Experimental design. Eight different fertilization treatments with three replicates were established: CK (control, without fertilization); NK (application of N + K); NP (application of N + P); PK (application of P + K); NPK (application of N + P + K), and another three different treatments combined with dairy FYM (M, NPKM, and 1.5NPKM). The details are provided in Table 1. The area of each plot was set as 30 m² (5 m × 6 m). Sweet sorghum was planted at a density of approximately 8–9 plants per 1 m², with a spacing of 60 cm between rows and with a spacing of 20 cm between plants.

Sample analysis. Soil samples (0–20 cm depth) were collected in September after harvesting every year. In each plot, the soil was collected from ten points randomly and mixed into one sample. After carefully removing the surface organic materials and fine roots, the soil samples were air-dried and sieved using a 2-mm sieve before determining the soil chemical properties. Soil pH_{H2O} and soil organic carbon (C_{tot}) were determined by the electrometric

instrument (Mettler Toledo FE20, Shanghai, China) and the dichromate wet oxidation method, respectively. Soil total salt (S_{tot}) content was estimated by weighing the evaporation residue of the soil/water (1: 5) extraction previously filtered on quantitative paper.

Table 1. The annual fertilization rate and fertilizer type of each treatment

Treatment	Nitrogen		Phosphorus		Potassium Manure		
	basal	topdressing	basal	topdressing (t/ha/year)	basal	topdressing	basal
CK	0	0	0	0	0	0	0
NK	108	72	0	0	27	27	0
NP	108	72	0	0	72	72	0
PK	0	0	27	27	72	72	0
NPK	108	72.0	27	27	72	72	0
M	0	0	0	0	0	0	12
NPKM	108	72.0	27	27	72	72	12
1.5NPKM	108	72.0	27	27	72	72	18

Statistical analysis. Statistical analyses were performed using Excel 2013 (Microsoft, Redmond, USA) and SPSS 25.0 software (IBM, Armonk, USA). Statistically significant differences were identified using a one-way ANOVA with the least significant difference (*LSD*) tests at the 0.05 level of significance. Data obtained from the triplicate measurements are presented as the mean \pm standard deviation (SD).

Weather data. Annual precipitation and temperature for Xinjiang were obtained at the China Meteorological Administration (<http://data.cma.cn/>).

Results and discussion

Annual average yield and stalk T. Biomass yields of the treated plots were significantly higher than that of CK (Figure 1A). The treatments NP, PK, NK, NPK and NPKM significantly increased the average biomass yields by 30–48%, without the marked differences between the years. Compared to CK, the average biomass of NPKM was the highest (94.06 t/ha), increased by 48.79%, but the average biomass yields of NP, PK and NPK were not significantly different from that of NPKM ($P > 0.05$). A significant difference ($P < 0.05$) in the average biomass yield among NPKM, 1.5NPKM, and M implies that 12 t/ha/year is the proper amount of manure for the combination with inorganic fertilizers to harvest higher yield. TSS is the weight percent of water-soluble solids in stem and is widely used as an approximation for with total sugar content. The TSS of sweet sorghum stems is shown in Figure 1B. The NPKM and M increased the TSS by 7.9% and 13.4% over CK, respectively, whereas no significant differences ($P > 0.05$) were observed among the average TSS for CK, PK, NK, NPK, 1.5NPKM and NP. The treatments NPKM and M increased the sugar TSS by 7.89% and 13.42%, respectively.

In conclusion, the 12 t/ha/year application rate of manure was recommended because NPKM emerged as the most efficient treatment in increasing the yield and TSS simultaneously.

The tendency of annual biomass and TSS. To get a more comprehensive understanding, the changes of biomass and TSS were analyzed over the study years. The biomass yields exhibited diverse tendencies in response to the different fertilization and varying weather course (Figure 2A). The biomass of CK significantly decreased with years, while the biomass of other treatments fluctuated over the years without showing a clear increasing trend.

The yield of CK significantly decreased by 40.66%, from 80.71 t/ha to 47.89 t/ha ($P <$

0.05). The biomass for the inorganically fertilised treatments (NP, NK, PK, and NPK) was extraordinarily high in 2010 and declined to a stable level in the following years. However, the biomass for the treatments with FYM and inorganics changed slightly. The biomass for NPKM kept the highest values during the later years. The changing of the TSS illustrated the same trend for all treatments over the years (Figure 2B). It increased from 2008 to 2009 and then decreased until 2018. At the end of the experiment, the highest TSS content was achieved with M, followed by NPKM and 1.5NPKM with slight differences. This result indicates that some nutrient sources from FYM can increase the TSS. Soil pH H₂O and Stot. Figure 3 represents the change in soil pH H₂O and Stot during 2008–2018. Soil pH H₂O for the treatments with FYM exhibited a steady increasing trend at the beginning of the experiment, whereas a quick decrease was observed for the other treatments. The pH H₂O for all treatments decreased sharply from 2012 to 2015 and then began to increase again (Figure 3A).

In 2018, the pH H₂O for CK was highest, with slight differences with the other treatments, and the pH H₂O for all treatments stabilized at 7.8–8.2. Treatment M decreased the soil pH H₂O value to a noticeable level. As illustrated in Figure 3B, the Stot was affected by different fertilization; first, the Stot for all treatments showed an upward trend, then declined dramatically, followed by a relatively stable level, and a similar trend was also observed for CK. After being stabilized, no significant differences across the treatments were observed in Stot; the Stot for treatment M was the lowest and reached a clear stabilized level. The results provided favorable evidence for preventing soil salinization and alkalization with a single FYM application.

Soil nutrient. The quantity of N, P and K distributed in each kilogram of topsoil was calculated according to the natural soil bulk density of this region 1.45 g/cm (Wang et al. 2016), the calculated inorganic N, P and K per kilogram of soil were 0.062 g/kg, 0.049 g/kg and 0.019 g/kg, respectively. The amounts of distributed nutrients in each kilogram of topsoil from 12 t/ha of FYM were 0.110g/kgN, 0.033 g/kg P, and 0.075 g/kg K, respectively.

Soil organic carbon. C_{tot} for the treatments ranged from 10.00 to 22.00 g/kg with years (Figure 4A). The C_{tot} for CK decreased gradually, whereas the C_{tot} for other treatments showed a wavy increasing tendency. The differences in C_{tot} between the CK and other treatments became more significant with years. At the end of the trial, the C_{tot} for 1.5NPKM was the highest, followed by NPKM with a slight difference. 1.5NPKM and NPKM increased the soil organic carbon by 46.25% and 44.23%, respectively. The C_{tot} for the treatments only with inorganic fertilisers also increased due to root biomass left in the soil.

Available potassium. The soil AK contents for some treatments, except for CK, NP and PK, increased steadily for the first 7 years, then decreased to stabilised level in 2016, and maintained till the end of the trial (Figure 4B).

Available nitrogen. The AN for all treatments increased in the early years of this trial (Figure 4C); CK had the minimum AN during this period. Then AN for all treatments decreased drastically until the end of the study period.

Available phosphorus. The annual variation in soil AP differed for the different treatments (Figure 4D). The AP for all treatments showed a steady increase for the first several years. Then only the two treatments M and NPKM, maintained the increasing trend while AP for the other treatments began to decline. Interestingly, AP for PK and NP decreased to relatively low values, although they were applied with phosphorus every year.

Weather impact. The annual precipitation, air temperature and their values for the growing season over the study period were plotted in Figure 5. Annual precipitation and precipitation in the growing season in 2010 and 2016 were higher. The more water supply in 2010 might give rise to the biomass yield for the inorganically fertilized treatments (Figure 2A) by elevating the water solubility of inorganic fertilizers; the effect of precipitation on yield was not further confirmed due to the absence of biomass data in 2016. No significant trend was revealed for the annual air temperature and temperature in the growing season.

Discussion

Soil $\text{pH}_{\text{H}_2\text{O}}$ affects nutrient availability to plants. Some components in the soil buffer against the $\text{pH}_{\text{H}_2\text{O}}$ changes caused by natural and anthropogenic inputs of acidic and basic fertilizers. Long-term fertilization of soil and cultivation of different crops may result in different consequences in different regions. In a cotton-chickpea cropping system in India's Vertisols, FYM combined with inorganic fertiliser markedly lowered the soil pH . Organic manure application contributed to the soil organic acids and caused a reduction in soil $\text{pH}_{\text{H}_2\text{O}}$ due to humic and fulvic acids from the microbial decomposition of organic manure, releasing of CO_2 in the soil (Walker et al. 2004) and to the nitrification of NH_4^+ (Chang et al. 1991). The phenolic groups in the soil also buffer $\text{pH}_{\text{H}_2\text{O}}$ at $\text{pH} > 8.5$ (Huang et al. 2011). In this study, the pH for all treatments undulated during the trial, and treatment only applied organic manure had the lowest soil $\text{pH}_{\text{H}_2\text{O}}$ at the end of this trial. The decrease in $\text{pH}_{\text{H}_2\text{O}}$ for the manured soil was in close agreement with the above findings. The role of FYM in reducing soil $\text{pH}_{\text{H}_2\text{O}}$ was clearly demonstrated after 11 years, possibly because of the slowness of organic manure decomposition in this arid experimental region.

Soil total salt. The topsoil salinity is highly variable, and the correlation between the salt concentration and the soil $\text{pH}_{\text{H}_2\text{O}}$ varies for the soils. In this study, the similar changing tendency of S_{tot} for all treatments, including CK, presented that S_{tot} was not affected by fertilization and sweet sorghum cultivation. Cotton cultivation without fertilization for 16 years resulted in a steady decrease in $\text{pH}_{\text{H}_2\text{O}}$ and salinity in this experimental region. The wavy fluctuation in S_{tot} was not in good agreement with the abovementioned result. The S_{tot} can be leached to below ground along with the irrigation and rainfall. Compared with the strong soil salt deposit caused by strong evaporation, the effects of irrigation, fertilization and cultivation on S_{tot} were relatively weak in this experiment. However, in 2016, the decrease in S_{tot} was probably caused by the relatively higher precipitation. More irrigation amount of water may be needed to overcome the strong evaporation.

Soil organic carbon. As reported by Mapfumo et al. (2007), C_{tot} can be increased with the application of extra organic fertilisation. However, the C_{tot} varies for the different soils mainly due to the decomposition rate of the organic matter. García-Ruiz et al. (2012) showed that the application of organic fertiliser resulted in no remarkable changes in C_{tot} for a short period due to the slow rate of the decomposition of organic matter. In the case of long-term organic fertilisation. Kouyaté et al. (2000) found that the effects of manure and crop residue on yield and soil organic carbon reached a significant level when sorghum was continuously cultivated for 8 years. In this experiment, significant differences in C_{tot} were observed for all fertilisation treatments at the end of the trial in comparison with that of the CK. The increasing C_{tot} for the treatments only with inorganic fertilisers might be attributed to the residual effect

caused by the slow humification of roots and plant residue.

Biomass and stem TSS. Soil nutrients need to be available for the plants for greater yield. All treatments increased the yield over the unfertilised control without showing an increasing trend over the years. The insignificant differences between the initial and final yields indicated that there are potentials for increasing the yields. The high sugar content of the sorghum stem is an important factor in assessing sorghum quality. The TSS for the treatments in this trial ranged from 14.87% to 22.92%, but the highest TSS was not obtained with the highest biomass yield. The treatment M accounted for the greatest TSS, followed by NPKM with a slight difference. The effect of fertilisers on the sweet sorghum sugar content varies with the environment and gene type, e.g., Holou and Stevens (2012) found that the application of nitrogen increased the sugar content in sweet sorghum. In contrast, another study (Almodares et al. 2009) observed that nitrogen's application decreased sugar content under the same conditions. In this trial, a relatively higher average value of TSS was observed in the treatments related to manure. The combination of extra manure with inorganics had a negative effect on the TSS in the case of 1.5NPKM, indicating that the amount of FYM in NPKM was proper for the soil type and cultivation of sweet sorghum in this experimental region.

Available potassium and nitrogen. The soil K^+ availability is highest above soil pH of 6.0, and the proper pH range for N availability is 6.0–9.0 (Weil and Brady 2016). The soil pH during the overall experimental period held above 7.4–8.6, which was at the interval of highly available K and N for plants. Thus, the gradual decrease in soil K and N was probably caused by the absorption by sorghum plants while the limited rainfall was not able to leach the soil N and K. The hardly increased biomass yields with years under decreasing available N and K condition requested for a more reasonable inorganic fertiliser application rate.

Available phosphorus. The phosphorus availability generally declines between 7.5 and 8.5 (Weil and Brady 2016). The pH for all treatments varied at 7.4–8.6 during 2008–2018, which indicated the less unavailability of phosphorus to a plant. The steady increase in available phosphorus in the previous years might be caused by the accumulation of P at this pH range. The reason for the subsequent decline in phosphorus for treatments, except for M and NPKM, needs further investigation.

References

1. Almodares A., Jafarinia M., Hadi M.R. The effects of nitrogen fertilizer on chemical compositions in corn and sweet sorghum // American-Eurasian Journal of Agricultural and Environmental Sciences. - 2009. - No 6. - Pp. 441-446.
2. Chang C, Sommerfeldt T.G., Entz T. Soil chemistry after eleven annual applications of cattle feedlot manure. // Journal of Environmental Quality. - 1991. - No. 20. - Pp. 475-480.

УДК 627.8.064.4

STUDING SCOUR HOLE DUE TO PLUNGING JETS

Alsadek Elias Sadek, Post-graduate student of the Department of Hydraulic Structures of Federal State Budgetary Educational Institution of Higher Education "Russian Timiryazev State Agrarian University", eliasalsadek@gmail.com